APPENDIX B SPECIAL-STATUS SPECIES

Note: This list was compiled in January 2002 based on information contained in the California Natural Diversity Database (CNNDB) and information from the United States Fish and Wildlife Service (USFWS).

Special Status Species				
Common Name	Reporting Source for Species Occurring in Area	Scientific Name	Status (Federal, State)	Habitat
		Bird	S	
Aleutian Canada goose	USFWS	Brant canadensis leucopareia	FD	Uses pastures and grain fields along the coasts of Oregon and California, and in California's Central Valley. Nest on maritime islands.
American peregrine falcon	USFWS	Falco peregrinus anatum	FD; CE	Wetlands, woodlands, forested areas, agricultural areas, and coastal habitats. Nesting sites on ledges.
Bank swallow	USFWS	Riparia riparia	СТ	Riparian, lacustrine, and coastal areas with vertical banks, bluffs, and cliffs with sandy soil. Nest in colonies in burrows dug into riverbanks.
Black tern	USFWS	Chlidonias niger	FSC; SC	Spring and summer in fresh emergent wetlands while breeding. Common on bays, salt ponds, river mouths and pelagic waters in spring and fall.
Burrowing owl	CNNDB/ USFWS	Athene cunicularia	SC, S2	Open grassland and desert habitats, in open parts of pinyon-juniper and ponderosa pine habitats. Uses rodent or other burrows for cover and nesting.
Cooper's hawk	GL-DEIR	Accipiter cooperii	SC	Oak woodlans, riparian or other forest habitat near

Special Status Species				
Common Name	Reporting Source for Species Occurring in Area	Scientific Name	Status (Federal, State)	Habitat
				water.
Ferruginous hawk	USFWS	Buteo regalis	FSC; SC	Open grasslands, sagebrush flats, desert scrub, low foothills surrounding valleys and fringes of pinyon-juniper habitats. Roosts in open area, usually in a lone tree or pole.
Golden eagle	GL-DEIR	Aquila chrysaetos	SC, Fully Protected	Rolling hills, mountain areas, sage-juniper flats and deserts.
Grasshopper sparrow	USFWS	Ammodramus savennarum	FSC	Tall and mixed grassland habitats including native prairies, hayfields, pastures, and fallow fields.
Greater sandhill crane	USFWS	Grus Canadensis tabida	СТ	Wet meadows. Tend to nest in open habitat or in the cover of bulrush and bur reed.
Lawrence's goldfinch	USFWS	Carduelis lawrencei	FSC	Open oak woodlands, mesquite, and riparian thickets.
Lewis' woodpecker	USFWS	Melanerpes lewis	FSC	Open pine-oak woodlands, coniferous forests, and riparian woodlands. Associated with burned and logged woodlands.
Little willow flycatcher	USFWS	Empidonax traillii brewsteri	CE	Wet meadows and montane riparian habitats with extensive willow thickets.
Loggerhead shrike	USFWS	Lanius ludovicianus	FSC; SC	Open habitats with scattered shrubs, trees, utility lines or other perches. Lowlands and foothills throughout California.
Long-billed curlew	USFWS	Numenius americanus	FSC; SC	Wet meadow habitat, Coastal estuaries, upland herbaceous areas, and croplands.

Special Status Species				
Common Name	Reporting Source for Species Occurring in Area	Scientific Name	Status (Federal, State)	Habitat
Mountain plover	USFWS	Charadrius montanus	FPT; SC	Short grasslands and plowed fields of the Central Valley.
Sharp- shinned hawk	GL-DEIR	Accipiter striatus	SC	Deciduous riparian forest at mid-elevation, conifer forest, and oak woodlands.
Short-eared owl	USFWS	Asio flammeus	FSC; SC	Grasslands, prairies, dunes, meadows, irrigated lands and saline and fresh emergent wetlands. Nests in depression in dry ground concealed in vegetation.
Swainson's hawk	CNNDB/ USFWS	Buteo swainsoni	СТ	Open desert, grassland, or cropland with scattered, large trees or small groves.
Tricolored blackbird	CNNDB/ USFWS	Agelaius tricolor	SC, S3	Emergent wetland vegetation with cattails, tules, and/or thickets.
Vaux's swift	USFWS	Cypseloides niger	FSC; SC	Redwood and Douglas-fir habitats with nests in large hollow trees and snags.
Western spadefoot	CNNDB	Scaphiopus hammondii	SC	Primarily in grassland habitats, also found in valley-foothill hardwood woodlands.
White-faced ibis	USFWS	Plegadis chihi	FSC; SC	Fresh emergent welands, shallow lacustrine waters, and the muddy ground or wet meadows and irrigated/flooded pastures and croplands.
White-tailed kite	CNNDB/ USFWS	Elanus leucurus	S3, Fully Protected	Lowland grasslands, agriculture, wetlands, oakwoodlands, savannah, and riparian habitats associated with open areas.

Special Status Species				
Common Name	Reporting Source for Species Occurring in Area	Scientific Name	Status (Federal, State)	Habitat
		Reptil	es	
California horned lizard	USFWS	Phrynosoma coronatum frontale	FSC; SC	Wide range of habitats from gravelly-sandy substrate containing scattered shrubs, to clearing in riparian woodlands.
Giant garter snake	USFWS	Thamnophis gigas	FT; CT	Marshes, sloughs, and slow-moving creeks, with nocturnal retreats in holes and mammal burrows.
Northwestern pond turtle	USFWS	Clemmys marmorata marmorata	FSC; SC	Pacific slope drainages from Washington to Baja California.
		Amphib	ians	
California red-legged frog	GL-DEIR	Rana aurora draytonii	FT; SC	Pools, ponds, slow streams, and marshes.
		Fish	1	
Central Valley fall/late fall- run Chinook salmon	USFWS	Oncorhynchus tshawytscha	FC; SC	The ocean and the Sacramento River and its tributaries.
Central Valley steelhead	USFWS	Oncorhynchus mykiss	FT	The ocean and the Sacramento and San Joaquin Rivers and their tributaries.
Green sturgeon	USFWS	Acipenser medirostris	FSC; SC	Pacific drainages, locally, the Sacramento-San Joaquin estuary.
Sacramento splittail	USFWS	Pogonichthys macrolepi- dotus	FT; SC	Primarily in the Sacramento-San Joaquin estuary.
Winter-run Chinook salmon	USFWS	Oncorhynchus tshawytscha	FE; CE	The ocean and the Sacramento River and its tributaries.

Special Status Species				
Common Name	Reporting Source for Species Occurring in Area	Scientific Name	Status (Federal, State)	Habitat
		Inverteb	rates	
California Linderiella fairy shrimp	CNNDB/ USFWS	Linderiella occidentalis	S2/S3	Seasonal pools in unplowed grasslands with old alluvial soils underlain by hardpan or in sandstone depressions.
Vernal Pool Tadpole Shrimp	CNNDB/ USFWS	Lepidurus packardi	FE; S2/S3	Vernal pools and swales in the Sacramento Valley containing clear to highly turbid water.
Valley Elderberry Longhorn Beetle	CNNDB/ USFWS	Desmocerus californicus dimorphus	FT; S2	Only occurs in the Central Valley of California in association with Blue Elderberry (Sambucus mexicana).
Vernal Pool Fairy Shrimp	CNNDB/ USFWS	Branchinecta lynchi	FT; S2/S3	In a static rain-filled pools in the central valley grasslands and central and south coastal mountains.
		Mamm	als	
Fringed myotis bat	USFWS	Myotis thysanodes	FSC	Roosts in caves, mines, and rock crevices within a variety of habitats.
Greater western mastiff-bat	USFWS	Eumops perotis californicus	FSC; SC	Open, semi-arid to arid habitats, including conifer and deciduous woodlands, annual and perennial grasslands, chaparral, and urban.
Long-eared myotis bat	USFWS	Myotis evotis	FSC	Woodland and forest habitats, roosting in rock crevices, under bark, and tree snags.
Long-legged myotis bat	USFWS	Myotis volans	FSC	Woodlands and forest habitats generally over 4,000 feet. Roosts in rock crevices, under bark, in tree snags, and cliffs.
Pacific	USFWS	Corynorthis	FSC; SC	All but alpine and sub-alpine

Special Status S	Species			
Common Name	Reporting Source for Species Occurring in Area	Scientific Name	Status (Federal, State)	Habitat
western big- eared bat		townsendii townsendii		habitats.
San Joaquin pocket mouse	USFWS	Perognathus inornatus	FSC	Dry, open grasslands or scrub area on fine textured soils in the Central and Salinas valleys.
Small-footed myotis bat	USFWS	Myotis ciliolabrum	FSC	Occurs in a variety of habitats, roosting in caves, crevices, and buildings.
Spotted bat	USFWS	Euderma maculatum	FSC	Arid or ponderosa pine forests, and marshlands. Roosts in small cracks in cliffs and stony outcrops.
Yuma myotis bat	USFWS	Myotis yumanensis	FSC	Variety of habitats from juniper and riparian woodlands to desert regions near open water. Associates with water and roosts in caves, attics, under bridges, mines, and similar places.
		Habita	ats	•
Alkali Meadow	CNNDB	-	S2	
Alkali Seep	CNNDB	-	S2	
Northern Hardpan Vernal Pool	CNNDB	-	S3	
Northern Volcanic Mud Flow Vernal Pool	CNNDB	-	S1	
Plants				
Big-scale Balsamroot	CNNDB	Balsamorhiza macrolepis var macrolepis	S2	Valley and foothill grassland, cismontane woodland.
Boggs Lake Hedge-	CNNDB/ USFWS	Gratiola heterosepala	CE, S3	Clay soils in marshes, swamps and vernal pools.

Special Status	Special Status Species				
Common Name	Reporting Source for Species Occurring in Area	Scientific Name	Status (Federal, State)	Habitat	
hyssop					
Dwarf Downingia	CNNDB	Downingia pusilla	S3	Valley and foothill grassland and several types of vernal pools.	
Hispid Bird's-Beak	CNNDB/ USFWS	Cordylanthus mollis ssp hispidus	FSC; S2	In damp alkaline soils in meadows, playas, and valley and foothill grassland.	
Legenere	CNNDB/ USFWS	Legenere limosa	FSC; S2	In beds of vernal pools.	
Red Bluff Dwarf Rush	CNNDB	Juncus leiospermus var leiospermus	S2	Chaparral, valley and foothill grassland, cismontane woodlands, and vernal pools.	

Notes:

Federal Status: FE=Federally Listed Endangered, FT=Federally Listed Threatened, FPE=Proposed for Federal Listing as Endangered, FPT=Proposed for Federal Listing as Threatened, FC=Candidate for Federal Listing, FSC=Federal Species of Concern, FD=Federal De-listed

State Status: CE=California Listed Endangered, CT=California Listed Threatened, CR=California Listed Rare, CC=Candidate for California Listing, SC=CDFG Species of Special Concern

NDDB Element Ranks: S1=Less than 6 element occurrences (EOs) OR less than 1,000 individuals OR less than 2,000 acres, S2=6-20 EOs OR 1,000-3,000 individuals OR 2,000-10,000 acres, S3=21-100 EOs OR 3,000-10,000 individuals OR 10,000-50,000 acres, S4=Apparently secure within California. NO THREAT RANK, S5=Demonstrably secure to ineradicable in California. NO THREAT RANK

Source: Quad Knopf, 2002; Granite Lakes Estates Draft EIR, EIP, 2001.